

TUBAL PREGNANCY AFTER ABDOMINAL BILATERAL TUBAL LIGATION

(A Case Report)

by

(Mrs) SHANTA JHA,* M.S. (Pat)

and

(Miss) MANJUBALA SHRIVASTAVA,** M.S. (Pat)

Introduction

A case of tubal pregnancy after tube ligation is reported.

Case Report

M. Kaur, 35 years, Hindu female, came to Rukmini Nurshing Home, Patna City on 15th February, 1981 at 4 P.M. for severe pain in abdomen and slight vaginal bleeding since morning of the same day. She had amenorrhoea for last one and a half months. Suddenly in the morning she developed acute pain in abdomen with fainting attack following which vaginal

bleeding also started. Her previous menstrual cycles were 4 to 5/40 days, regular. Her last menstrual period was 2nd January, 1981.

She had 6 full term normal deliveries. Past history nothing was particular except that she had abdominal bilateral Tube ligation done 5 years back.

She was pale, markedly cyanosed. Pulse rate 120/mt., regular, low volume. B.P. was 90/60 mm. of Hg.

There was tenderness in lower abdomen. On vaginal examination, uterus was found to be retroverted and normal in size, right fornix was clear. There was a boggy, tender mass in the left fornix. Cervical excitation test was positive. Rupture of tubal pregnancy was suspected. An urgent haemoglobin estimation was done, it was 60 per cent.

Laparotomy was done under general anaesthesia. Left fallopian tube at the tubectomy site towards the fimbrial end was found ruptured. Left sided salpingectomy was done.

* Tutor, Deptt. of Preventive & Social Medicine, Patna Medical College, Patna.

Medical Officer Post-partum Programme, Deptt. of Obstetrics and Gynaecology, Patna Medical College Hospital, Patna.

Accepted for publication on 1-10-1981.