

# COLLISION TUMOUR OF CORPUS UTERI WITH CO-EXISTING BRENNER TUMOUR—A HISTOLOGICAL CURIOSITY

by

S. RASTOGI,\*  
V. K. PRATAP,\*\*

and

VINITA DAS,\*\*\*

Brenner tumour associated with endometrial carcinoma, leiomyosarcoma and cervical dysplasia is a histological rarity. This case is being presented due to its varied histological heterogeneity.

## CASE REPORT

P.W. aged 55 years, H.F. P<sub>3</sub> + O, postmenopausal for 10 years was admitted in the department of Obstetrics and Gynaecology, L.L.R.M. Medical College, Meerut on 29-10-1981 for postmenopausal bleeding for last 10 months.

Obstetric History—P<sub>3</sub> + O. Last delivery was 25 years back.

Uterus enlarged to 6 weeks size, firm, mobile, left fornix—clear, right fornix a solid mass 3" x 4" with well defined margins, mobile, non-tender was felt.

Provisionally a diagnosis of ? Cancer body

\*Reader.

\*\*Reader.

\*\*\*Lecturer.

Departments of Obstetrics and Gynaecology and Pathology, LLRM Medical College, Meerut.  
Accepted for publication on 9-9-82.

uterus with right sided ovarian tumour was made.

## Treatment:

Abdominal panhysterectomy with removal of cuff of vagina was done on 10-11-1981. Patient made uneventful recovery and discharged on 19-11-1981 in satisfactory condition and upto now the patient is perfectly well.

## Histopathology:

Right sided Brenner tumour with endometrial carcinoma with leiomyosarcoma with cervical dysplasia.

## Discussion

An association of Brenner tumour with endometrial cancer and with adenomatous hyperplasia of endometrium has been reported by Ming and Goldman (1962).

## Reference

1. Ming, S. and Goldman, H.: Am. J. Obstet. Gynec. 83: 666, 1962.

---

See Figs. on Art Paper VI