

ECTOPIC BREAST ON VULVA

(A Case Report)

by

(Mrs.) C. D. DOIFODE

and

(Miss) L. K. BICHILE

Abberent breast tissue may be recognised on the vulva since this area is in the nipple line. The gross appearance is usually suggestive of breast due to presence of nipple like appearance of slightly pigmented lesion. Lactation may also be recognised in most of the cases after delivery.

CASE REPORT

Mrs. P.M. was admitted to Medical College Hospital, Aurangabad on 17th August, 1983 for swelling over vulva since 8 years. To begin with it was to small size and it gradually increased to present size. No history of increase in

From: Govt. Medical College, Aurangabad-431 001.

Accepted for publication on 12-12-1983.

size after delivery. She had 6 full term deliveries and last child birth was 1 year.

On examination of vulva there was pedunculated growth arising from left labia majora about 3" x 2" x 1" firm nontender. There was ulceration over dependent area. No pigmentation or elevated area suggestive of nipple areola. Speculum and vaginum examination did not reveal anything abnormal.

Tumour was excised under spina anaesthesia. There was no difficulty in excision. Her post-operative period is uneventful.

Histopathology Report

Slide shows stratified squamous epithelial lining with hairs follicles and sebacious gland. Underneath breast tissue composed of groups of acini and ducts separated by fibrous stroma suggestive of breast (Fig. 1).

See Fig. on Art Paper IX