

## A case of Carcinoma Cervix Occurring in a Patient with Uterovaginal Prolapse.

K. C. Dev, Nandini Gupta, Anup Bhuniya, Polly Saha, Abhinebesh Chatterjee

Eden Hospital, Department of Obstetrics & Gynaecology, Medical College, Calcutta, 88, College Street, Calcutta-700 073.

### Introduction:

Many patients come to our O. P. D. with procidentia and decubitus ulcer which they have neglected for years. It is rare to find a case of cervical cancer arising in an old decubitus ulcer. This rare complication is being reported here.

A 65 years old patient P8+0, last delivery 21 years ago, Menopause 14 years ago presented in the O.P.D. with H/O something coming down per vaginum for last 15 years and recent onset of post menopausal bleeding.

Examination revealed procidentia with an ulcerative growth arising from the cervix. The parametrial involvement was upto the lateral pelvic walls.

### Investigation

- Hb% 6.5%
- Urea, creatinine, PPBS-CXR, Urine,
- Stool = NAD
- Cx. Biopsy- infiltrating Keratinising squamous cell cancer.

### Management:

The patient received 5 units of blood transfusion to raise Hb to 11g%. Regular dressing of the decubitus ulcer was done.

Patient was referred to the Radiotherapy unit as it was judged to be inoperable. She was given telecesium therapy of Dose 4 SGY over a period of 4 weeks.



Fig 1