


Genital Prolapse with Chemical Vesico Vaginal Fistula

Sebanti Goswami, Debjani Bhadra, Mamata Pradhan

Department of Gynaecology & Obstetrics, Calcutta National Medical College & Hospital, Calcutta

Genital prolapse with chemical VVF is very rare. This case is presented for its rarity. Mrs. T.B. 43 yrs., P7+0 (LI-6; P7 → Caesarean section); attended the GOPD of Calcutta National Medical College and Hospital on 30.11.99 with the complaints of something coming down P/V for last 3 years along with continuous dribbling of urine for last one month and irregular menstruation for 1 year. She had gone to a quack 3 months back who prescribed a medicine to be applied on the prolapsed site for its remedy, which she did for a month. Suddenly one day she felt the continuous dribbling of urine per vagina. O/E – General examination within normal limits; P/A – NAD; On inspection – 2nd degree uterine prolapse with moderate degree cystocele with a 3cm x 3cm vesico-

vaginal fistula with prolapse of the bladder mucosa detected (midvaginal fistula) (Photograph) Rectocele – mild. P/V – ut. NS; Fornices – clear; Palpation of the fistula showed no significant fibrosis around the fistula margin. Routine investigation → within normal limits; Cystoscopy – Fistula just proximal to the bladder neck; IVP – mild hydronephrotic change in the right ureter; Endometrial biopsy – proliferative endometrium. Vaginal hysterectomy with PFR alongwith repair of the fistula done under GA. Post operative period was uneventful. Patient was discharged from the hospital on the 16th post operative day. Follow up examination after 6 weeks revealed no complaints.



Photograph I