

Transvaginal Sacrospinous Colpopexy for Vault Prolapse and for Marked Uterovaginal Prolapse

N Rajamaheswari, Gunasekaran Karthik

Govt. Kasturba Gandhi Hospital, Madras Medical College, Chennai-600005.

OBJECTIVE S – To assess the efficacy of transvaginal sacrospinous colpopexy in the treatment of vault prolapse and marked uterovaginal prolapse. **METHODS** – A retrospective study of 30 patients who underwent sacrospinous colpopexy procedure from July 1999 to June 2000 was done. Of these, 10 had vault prolapse and 20 had marked uterovaginal prolapse. Bilateral sacrospinous colpopexy was done in 14 patients and 16 underwent the procedure unilaterally. Enterocele repair, posterior vaginal repair and anterior vaginal repair were done when indicated and 20 with uterovaginal prolapse underwent vaginal hysterectomy. All patients were reviewed postoperatively at the end of 1 month and 3 months, and half yearly thereafter. **RESULTS** – Duration of follow up ranged from 2¹/₂ to 3¹/₂ years. None of the patients developed recurrence of prolapse in the anterior compartment, the posterior compartment or in the vault. Complications were limited to transient lower limb weakness in two patients, both of whom recovered uneventfully. **CONCLUSION** – Sacrospinous colpopexy is a simple and effective procedure in the treatment of vault prolapse and marked uterovaginal prolapse. It is a procedure that should be learnt and applied more frequently considering its many advantages.

Key words : colpopexy, sacrospinous colpopexy, vault prolapse, uterovaginal prolapse

Introduction

With the dawn of the new century, the problem of pelvic organ prolapse has been receiving tremendous attention. An increase in longevity and a yearning for an active lifestyle with maintenance of coital function has made women keen for permanent solutions for pelvic organ prolapse. The aim of any prolapse surgery is to correct the anatomical problem, relieve symptoms and restore function. The surgical procedures aimed at effective vault support have been many. The transvaginal sacrospinous colpopexy (SSCP) is one such procedure which has become increasingly popular. The first attempts by Zweife¹ in 1892 were followed by Richter² in 1968 and later were popularized by Randall and Nichols³ in 1971 in the USA.

The procedure involves fixing the vaginal vault to the sacrospinous ligament with permanent or delayed absorbable suture material. Indications for sacrospinous colpopexy include symptomatic prolapse of vaginal vault, nulliparous prolapse needing preservation of fertility and prophylaxis against vault eversion at the time of vaginal hysterectomy.

Material and Methods

Thirty multiparous women having genital prolapse

Paper received on 01/04/03 ; accepted on 21/09/03

Correspondence :

Dr. N. Rajamaheswari

Department of Urology / Urogynecology

Govt. Kasturba Gandhi Hospital, Madras Medical College, Chennai - 600 005.

underwent SSCP from July 1999 to June 2000. They were evaluated with a standardized protocol based on a) history b) gynecological and urological examinations c) cystoscopy d) stress test with the patient in standing position after reduction of the prolapse with pessary, and e) urodynamic studies (cystometry and pressure flow) as and when needed. Table I gives details of presenting complaints and associated pelvic floor defects seen in patients. A good knowledge of the working anatomy of the sacrospinous ligament and its surrounding structures is required for minimizing complications during surgery and achieving maximum results. The sacrospinous ligament is attached medially by a broad base to the side of the lower part of sacrum and upper part of coccyx. It narrows as it passes laterally, where its apex is attached to the ischial spine. Randall and Nichols³ described the sacrospinous ligament as a cord like structure lying in the substance of the coccygeus muscle. In fact they are so similar as to be referred to as the coccygeus sacrospinous ligament (C-SSL). Verdeja et al⁴, on dissecting 24 human cadavers noted that the pudendal complex and sciatic nerve were found 0.90 to 3.30 cms medial to the spine, making them prone to injury.

Surgical procedure

The SSCP begins with a longitudinal incision in the posterior vaginal wall from introitus to the vault. The rectovaginal space is defined. Any existing enterocele is dissected, opened and high ligation of the sac performed. By breaking the rectal pillar, the ischial spine is felt. A long needle holder with an atraumatic prolene suture

No. '1' is used to take a bite on the sacrospinous ligament at a point 3 cms medial to the spine. A second bite is taken using the same suture 1 cm medial to the first. Both sutures are loaded on to needles and brought out through the vaginal wall at the level of the vault. The anterior and posterior defects if present are then corrected. The sacrospinous sutures are finally tied taking the vault to the ligament.

Results

All patients were operated under regional anesthesia. The details of concomitant surgical procedures done are given in Table II. Operating time ranged between 60 and 90 minutes, when vaginal hysterectomy was included, and 30 and between 60 minutes when only repair of vault prolapse with scarospinous colpopexy was done. Cystoscopic removal of vesical stone was done in one

patient. One patient underwent repair of prolapsed urethral mucosa. Both these procedures were done prior to the primary procedure. Two patients with stress urinary incontinence underwent surgery with tension free vaginal tape as described by Ulmsten et al⁵ after the primary procedure. There was no need for any blood transfusion. Urinary catheter was left in for 3 days postoperatively in all cases. Postoperative complication was limited to transient bilateral lower limb weakness in two patients which recovered in 4 or 5 days time. All patients were discharged in 3 to 5 days. The patients who underwent TVT were completely cured of SUI. Follow-up ranged from 2 1/2 years to 3 1/2 years. The patients were examined after discharge at 1 month and 3 months, and thereafter at 6 monthly intervals and were found to be symptom free without any recurrence during the follow up period ranging from 2 1/2 to 3 1/2 years.

Table I : Associated conditions and pelvic floor defects.

	Cystocele	Rectocele	Enterocele	Stress urinary incontinence	Bladder Stone	Urethral mucosal prolapse
Uterovaginal prolapse (n=20)	20	15	10	2	1	1
Vault prolapse (n=10)	10	8	10	-	-	-

Table II : Concomitant procedures done along with sacrospinous colpopexy

	Sacrospinous colpopexy		Concomitant procedures						
	Bilateral	Unilateal	VH	AC	PC	ER	TVT	Bladder stone removal	Urethral mucosal repair
UVP	20	8	20	20	15	10	2	1	1
VP	10	10	0	-	10	8	10	-	-

VP – Vault Prolapse
 AC – Anterior vaginal repair
 TVT - Tension free vaginal tape
 UVP – Uterovaginal prolapse
 PC – Posterior vaginal repair
 VH – Vaginal hysterectomy
 ER – Enterocele repair.

Discussion

Pelvic organ prolapse can be the result of chronic increase in abdominal pressure exerted on normal pelvic organs or defective support responding inadequately to normal intraabdominal pressure. Incidence of massive vaginal vault prolapse following hysterectomy is 2 to 3.6 / 1000 women years⁶. It is however interesting to note that vault prolapse rates reported after hysterectomy for uterine

prolapse, were higher viz. 15/1000 women years⁶. Marked vault or uterovaginal prolapse requires procedures that provide adequate vaginal depth and prevent recurrence^{7,8}. The apex of the vagina lies almost horizontal over the levator plate. The sacrospinous colpopexy⁹ procedure re-establishes the upper vagina over the levator plate. The patients in our study underwent not only SSCP but also concomitant repair of rectocele, cystocele and enterocele when required.⁴

Associated stress urinary incontinence was also corrected. The reported incidence of postoperative stress urinary incontinence after correction of uterovaginal prolapse is 15-80%¹⁰. In all our patients the prolapse was reduced with pessary before checking for stress urinary incontinence by cough test. Those who tested positive underwent urodynamic studies. The TVT procedure was done in patients having genuine stress incontinence as it was a highly effective minimally invasive procedure with least morbidity. The technique itself varies from center to center with surgeons using Deschamps¹¹ ligature carrier or Miya Hook¹² and / or the endostitch. We combined anterior colporrhaphy in all our patients and did not come across recurrent cystoceles in the follow up. As the sacrospinous colpopexy retroverts the vault, patients are highly prone to anterior compartment defects¹³.

Even though there is controversy as to whether there is any advantage of the bilateral procedure over the unilateral one, we preferred the bilateral one in patients with vault prolapse. Also a redundantly wide vault was better dealt with by bilateral sacrospinous colpopexy. None of the patients complained of dyspareunia.

In obese women with deep pelvis and less prominent ischial spines, identifications of sacrospinous ligament may be difficult. If the stitch does not include the sacrospinous ligament, it may cut through while it is anchored to the vagina.

During surgery, rectum, pudendal nerve and pudental vessels can get injured and due care should be taken to avoid such injuries. This surgery is contraindicated if the patient had rectal surgery or drainage of pelvic abscess in the past.

References

- Zweifel P. Vorlesungen uber Klinische Gynecologie. Berlin : Hirschwald 1982:407 - 15.
- Richter K. Die Chirurgische anatomie der vaginaefixatio sacrospinous vaginalis. Ein Beitrag zur operativen Behandlung des Scheidenblindsach prolapses. *Geburtshilfe Frauenheilkd* 1968;78:321-7.
- Randall CL, Nichols DH. Surgical treatment of vaginal eversion. *Obstet Gynecol* 1971;38:327-32.
- Verdeja AM, Elkins TE, Odoi A et al. Transvaginal sacrospinous colpopexy: anatomic landmarks to be aware of to minimize complications. *Am J Obstet Gynecol* 1995;173:1468-9.
- Ulmsten U, Henriksson L, Johnson P et al. An ambulatory surgical procedure under local anesthesia for treatment of female urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunction* 1996;7:81-6.
- Mant J, Painter R, Vessey M. Epidemiology of genital prolapse: observations from the Oxford Family Planning Association study. *Br J Obstet Gynecol* 1997;104:57-85.
- Addison WA, Livengood CH, Sutton GP et al. Abdominal sacral colpopexy with Mersiline mesh in the retroperitoneal position in the management of post-hysterectomy vaginal prolapse and enterocele. *Am J Obstet Gynecol* 1985;153:140-6.
- Maloney JC, Dunton CJ, Smith K. Repair of vaginal vault prolapse with abdominal sacropexy. *J Reprod Med* 1990;35:6-10.
- Morley GN, DeLancey JO. Sacrospinous ligament fixation for eversion of the vagina. *Am J Obstet Gynecol* 1988;158:872-9.
- Bump RC, Hurt GW, Theofrastous JP et al. Randomised prospective comparison of needle colposuspension versus endopelvic fascia plication for potential stress incontinence prophylaxis in women undergoing vaginal reconstruction for stage III or IV pelvic organ prolapse. *Am J Obstet Gynecol* 1996;175:326-35.
- Nichols DH. Sacrospinous fixation for massive eversion of the vagina. *Am J Obstet Gynecol* 1982;142:901-4.
- Miyazaki FS. Miya hook ligature carrier for sacrospinous ligament suspension. *Obstet Gynecol* 1987;70:286-8.
- Holley RJ, Varner RE, Gleason BP et al. Recurrent pelvic support defects after sacrospinous ligament fixation for vaginal vault prolapse. *J Am Coll Surg* 1995;180:444-8.